

IV PRE-SURGICAL INSTRUCTIONS

- 1. Discontinue aspirin and any medications containing aspirin for 2 weeks prior to your surgical appointment, unless instructed by your physician <u>not</u> to discontinue this medication.
- 2. Discontinue Advil or any other non-steroidal anti-inflammatory medication one-week prior to your surgical appointment. You may substitute Tylenol for these medications if you are medically able to take Tylenol.
- 3. Discontinue anything herbal, Vitamin E, Ginseng, and Fish oil one week prior to your surgical appointment.
- 4. Inform this office if you are taking any other anticoagulant medication, i.e., Plavix, Coumadin, etc. ** You will need to be off these medications prior to surgery if your physician allows it.
- 5. Take any prescribed antibiotic as directed for heart murmur or Mitral Valve Prolapse.
- 6. Take all routine medications except for those listed above as directed by your physician. Please inform this office of all routine medications you are currently taking.
- 7. Please inform this office of all allergies to medications.
- 8. Plan to rest at home the remainder of the day and the following day after your surgical appointment. Plan to have a caretaker the remainder of the day and the following day.
- NOTHING TO EAT OR DRINK AFTER MIDNIGHT the night before surgery. DO take routine medications as directed but only with a small amount of water. If you were given special instructions during your consult please follow these instructions. These include:
- 10. Following surgery you should have cold liquids for the next 2 days. Protein shakes, Instant Breakfast, Ensure, ice cream, and yogurt are good choices. Straws are <u>NOT</u> to be used.
- 11. START taking a multi-vitamin and vitamin c no more than 1000mg today and continue until you have healed from your surgical procedure if you are not taking on a regular basis already. Multi-Vitamin must have zinc and magnesium.
- 12. Take all medications as prescribed pre op and post op.
- 13. Please have someone available to drive you to the office and remain until the surgery is over to take you home. Due to the sedation, you <u>CANNOT</u> drive for at least 24 hours after your surgery.